



Overview

Duloxetine (Cymbalta®) belongs to a group of medications called antidepressants. Duloxetine may be used to treat several conditions.

What is *duloxetine* used for?

Like many medications used to treat childhood disorders, duloxetine has not been approved by Health Canada for use in children and adolescents. When the potential benefits (e.g., reducing your symptoms) of using duloxetine outweigh the potential risks (e.g., the side effects), many doctors may prescribe it to treat:

- Depression
- Depression associated with body pain
- Generalized anxiety disorder
- Some Anxiety disorders such as panic disorder, separation anxiety disorder, obsessive-compulsive disorder (OCD) and posttraumatic stress disorder (PTSD)

Your doctor may be using this medication for another reason. If you are unclear why this medication is being prescribed, please ask your doctor.



How does *duloxetine* work?

Duloxetine is a “Serotonin and Norepinephrine Reuptake Inhibitor” (SNRI). This means it increases the amount of certain chemicals in the brain called serotonin and norepinephrine. It is believed that these brain chemicals are not working well in people who are depressed. It may also slightly increase another chemical in the brain called dopamine. The exact way that duloxetine improves the symptoms of depression is still not fully known.

How well does *duloxetine* work in children and adolescents?

In some studies of children and adolescents with depression, medications like duloxetine have been found to help about **5-6 people out of every 10 who are treated**. Duloxetine has been studied in youth with depression as well as youth with anxiety disorders. Some of these trials have found that duloxetine was not better at lowering the symptoms of depression than a placebo (an inactive pill that looks like the medication).

In general, depressed youth prescribed duloxetine for 2-3 months should notice an improvement in their depressive symptoms (*such as improved mood, better sleep, more energy, and improved concentration*). Whenever possible, the addition of behavioural therapy such as *Interpersonal Therapy (IPT) for depression* or *Cognitive Behaviour Therapy (CBT)* to this medication may help to increase the potential for benefits.

How should *duloxetine* be taken?

Duloxetine is usually taken once or twice a day with or without food (food may help reduce early symptoms of nausea). This medication should be taken at the same time(s) each day as directed by your doctor. Try to connect it with something you do each day (like eating breakfast or brushing your teeth) so that you don't forget. Try to avoid alcohol when taking duloxetine.

Swallow delayed release duloxetine capsules whole with fluid. The capsules and their contents should NOT be opened, chewed or crushed, nor should the contents be sprinkled on food or mixed with liquids.

Usually, your doctor will start with a low dose of duloxetine. This dose will be slowly increased based on how you respond to it. You and your doctor can then discuss the best dosage to stay on based on how you tolerate this medication (how well the medication is working and how you are doing with the side effects of the medication) and how well it helps to decrease your symptoms.

When will *duloxetine* start working?

Duloxetine must be taken for 3 to 6 weeks before you begin to feel better. Different symptoms may start to improve at different times. For example, improvements in sleep, appetite and energy may be seen within the first 2 weeks. Sometimes, others will notice improvements in you before you do. Full beneficial effects may take 4 to 8 weeks (or longer). **Since this medication takes time to work, do not increase, decrease or stop taking it without discussing with your doctor first.**

If you are not feeling better within 6 to 8 weeks, your doctor may recommend you take a different medication. There is a small chance that depression or anxiety symptoms may worsen or that you may experience increased thoughts of self harm during the first months of taking this medication (see section on side effects). If this happens, tell your doctor **IMMEDIATELY**.



How long do I have to take *duloxetine*?

This depends on the symptoms you have, how frequently they occur and how long you have had them. Most people need to take this medication for at least 6 months. This allows time for your symptoms to stabilize and for you to regain functioning. After this time, you and your doctor should discuss the benefits and risks of continuing treatment.

If you have had several episodes of severe depression and you tolerate this medication well, you may be asked to take this medication for an indefinite amount of time. By continuing to take this medication, you significantly decrease the chance that you may have another episode of depression. **Do NOT stop taking this medication (even if you are feeling better) without discussing it with your doctor first. If you stop taking this medication suddenly, it is possible that your symptoms may return or you may have a bad reaction.**

Once you have started taking this medication, you and your doctor will need to monitor for both the beneficial and unwanted effects. Your doctor will likely check your progress and discuss changes in symptoms during the next 3 months to confirm that this medication is working properly and that possible side effects are avoided. At this time, you can discuss how long you might need to take this medication.



Tip: Use the *Antidepressant Monitoring Form for Children and Adolescents* to help measure your progress on this medication.

Is *duloxetine* addictive?

No, duloxetine is not addictive. You will not have “cravings” for it like some people do with nicotine or street drugs. If you and your doctor decide it is best for you to stop using duloxetine, your doctor will explain how to safely lower the dose so you won’t feel any unpleasant “flu-like” effects (chills, nausea, vomiting, dizziness, tingling in hands and feet, muscle aches, fever and electrical sensations) as your body adjusts to being without it.

What are the side effects of *duloxetine* and what should I do if I get them?

As with most medications, side effects may occur when taking duloxetine. Most side effects are mild and temporary. Side effects may occur before any of the beneficial effects. It is possible for some individuals to experience side effects that they feel are serious or long lasting. If this occurs, speak to your doctor about ways to manage them. On the next page are some of the common side effects and potentially serious side effects of taking this medication. In brackets are suggested ways to lessen these effects.

Common side effects

Side effects may be more common when starting a medication or after a dose increase. If any of these side effects is too troublesome for you, please discuss them with your doctor, nurse or pharmacist.

- Constipation (*increase exercise, fluids, vegetables, fruits and fiber intake*)
- Decreased appetite (*try eating smaller, more frequent meals*)
- Drowsiness (*take the dose at bedtime; this usually lessens over time*)
- Difficulty sleeping (*try taking the medication earlier in the day*)
- Dizziness (*try getting up slowly from a sitting or lying down position*)
- Dry mouth (*try chewing sugarless gum, sour candies, ice chips, or popsicles*)
- Energized/agitated feelings (*avoid caffeine from energy drinks, colas and coffee*)
- Excessive sweating (strong antiperspirants can help; talk with your doctor or pharmacist)
- Headache (*try using a pain reliever like acetaminophen (plain Tylenol[®])*)
- Changes in sexual performance or interest (*discuss with your doctor*)
- Stomach aches or nausea (*try taking the duloxetine dose with food*)
- Unusually vivid dreams

Uncommon side effects (e.g., those that occur in less than 5% of patients)

Contact your doctor **IMMEDIATELY** if you have any of these side effects:

- Change in mood to an unusual state of excitement, irritability or happiness
- Muscle twitches or stiffness
- Seizures (also called fits or convulsions)
- Skin rash, itchy skin or hives
- Thoughts of self harm, hostility or suicide
- Uncomfortable sense of inner restlessness or agitation
- Unusual bruising or bleeding



What precautions should my doctor and I be aware of when taking *duloxetine*?

Tell your doctor or pharmacist if you:

- begin taking any other new medication (prescription or non-prescription), since several other medications can interact with duloxetine
- feel drowsy, dizzy or slowed down. Duloxetine can make some individuals experience these temporary side effects. Duloxetine may increase the effects of alcohol, resulting in more sedation or dizziness. if you feel this way, it is important to avoid operating heavy machinery or driving a car.
- have a history of diabetes, heart disease, thyroid disease, kidney or liver disease or seizures
- have any allergies or have experienced a reaction to a medication
- have any changes in mood or thoughts of self harm
- if you develop any new medical problem while you are taking an SSRI
- miss a period, become pregnant or are trying to become pregnant or are breast-feeding

What special instructions should I follow while using *duloxetine*?



- Keep all appointments with your doctor and the laboratory.
- Do not allow anyone else to use your medication.
- It is a good idea to have a visit or telephone call with your doctor within 1-2 weeks after you start taking duloxetine, and then periodically after that to see how well the medication is working, how well you are tolerating the medication, and to discuss any problems you may have.

What should I do if I forget to take a dose of *duloxetine*?

If you miss a dose of this medication, take it as soon as possible. However, if it is almost time for your next dose (e.g., within 4 hours), do not take the missed dose or double your next dose. Instead, continue with your regular dosing schedule.



What storage conditions are needed for *duloxetine*?

- Keep this medication in the original container, stored at room temperature away from moisture and heat (e.g., not in the bathroom or kitchen).
- Keep this medication out of reach and sight of children.

You may wish to share this information with your family members to help them to understand your treatment options. Since every person's needs are different, it is important that you follow the advice provided to you by your own doctor, nurse and/or pharmacist and speak to them if you have any questions about this medication.

Developed by the health care professionals of Child & Adolescent Mental Health Program and reviewed by the staff of the Kelty Resource Centre.